



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**Proteases.**—VINES,<sup>23</sup> in continuing his work on the proteolytic enzymes of plants, finds that both malt extract and taka-diastrase (Parke Davis and Co.) contain enzymes capable of digesting fibrin and of splitting peptone. From malt extract he has isolated the peptone-splitting enzyme free from the fibrin-digesting body, and from taka-diastrase he separated each from the other. Both these enzymes seem to act best in acid media. In animal tissues there are two fibrin-splitting enzymes:  $\alpha$  protease, weak and acting in basic media; and  $\beta$  protease, more powerful and acting in acid media. By special methods of preparation, VINES obtained a protease which acted best in neutral and basic media. This perhaps corresponds to the  $\alpha$  protease of animal tissues. The ereptases, peptone-splitting enzymes, of animal tissues act best in basic media. VINES'S work shows that plant ereptases act in acid media. As to terminology, one is inclined to believe that VINES could adopt profitably that of animal workers as given by VERNON.<sup>24</sup>—WILLIAM CROCKER.

**A new genus of Cordaitales.**—SCOTT and MASLEN<sup>25</sup> have described a new genus (*Mesoxylon*) of Cordaitales from the calcareous nodules of the Lower Coal-measures of Lancashire. It is intermediate between *Poroxylon* and *Cordaïtes*, as its name implies, including five species which have been referred heretofore to these two genera. The combination of characters is the anatomical habit of *Cordaïtes* and the centripetal xylem of *Poroxylon*. The pith is relatively large and discoid (as in *Cordaïtes*); the wood is dense, with narrow pith rays and relatively small tracheids; the leaf traces are double, but divide before entering the leaf; the centripetal xylem is present in the leaf traces at the margin of the pith (as in *Poroxylon*) and throughout their course to the leaves. The genus is thought "to completely bridge the gap, so far as anatomy is concerned, between the Poroxyleae and the Cordaïteae," and helps to connect the cordaïtean and later forms (excepting cycadophytes) with the "pteridosperms."—J. M. C.

**"Bars of Sanio" in Coniferales.**—The "bars of Sanio" are "folds" of cellulose to be observed in the walls of tracheids as horizontal or more or less semicircular markings, which stand out clearly with proper staining. Miss GERRY<sup>26</sup> has investigated their distribution among the Coniferales, and has discovered that they furnish a constant and useful character in the determination of fossil woods. They were found in 35 of the living genera, but do not occur in *Agathis* and *Araucaria*, nor in the mesozoic araucarians. Since they do occur in the podocarps, it is estimated that this group is more closely related to the Abietineae than to the Araucarineae, a conclusion which contradicts a growing conviction based

<sup>23</sup> VINES, S. H., Proteases of plants. *Annals of Botany* **24**:213-222. 1910.

<sup>24</sup> VERNON, H. M., Intracellular enzymes. London: John Murray. 1908.

<sup>25</sup> SCOTT, D. H., and MASLEN, A. J., On *Mesoxylon*, a new genus of Cordaitales (preliminary note). *Annals of Botany* **24**:236-239. 1910.

<sup>26</sup> GERRY, ELOISE, The distribution of the "bars of Sanio" in the Coniferales. *Annals of Botany* **24**:119-124. *pl.* 13. 1910.